

On a Rigorous Proof of the Joos-Zeh Formula for Decoherence in a Two-Body Problem ¹

Alessandro Teta

Dipartimento di Matematica Pura e Applicata, Università di L'Aquila, Italy
e-m: teta@univaq.it

Abstract

We consider a simple one dimensional system consisting of two particles interacting with a δ -potential and we discuss a rigorous derivation of the asymptotic wave function of the system in the limit of small mass ratio. We apply the result to the explicit computation of the decoherence effect induced by scattering in a concrete example of quantum evolution.

In this note we shall briefly discuss the asymptotic form of the wave function of a two-particle system in the limit of small mass ratio.

The interest in this problem is motivated by the analysis initiated by Joos and Zeh ([JZ]) of the mechanism of decoherence induced on a heavy particle by the scattering of light ones (see also [GF],[T],[BGJKS],[GJKKSZ],[D]).

The basic idea for the analysis of the process is that the small mass ratio produces a separation of two characteristic time scales, one slow for the dynamics of the heavy particle and the other fast for the light ones.

Following this line, in [JZ] the elementary scattering event between a light and a heavy particle is described by the instantaneous transition

¹Proceedings of the Conference “Multiscale Methods in Quantum Mechanics: Theory and Experiments”, Roma, December 16-21, 2002.

$$\phi(R)\chi(r) \rightarrow \phi(R)(S_R\chi)(r) \quad (1)$$

where ϕ and χ are the initial wave functions of the heavy and the light particle respectively and S_R is the scattering operator of the light particle when the heavy particle is considered fixed in its initial position R .

In (1) the initial state is chosen in the form of a product state, i.e. no correlation is assumed at time zero; moreover the final state is computed in a zero-th order adiabatic approximation for small values of the mass ratio.

Formula (1) gives a simple and clear description of the scattering event; nevertheless the approximation chosen is rather crude in the sense that time zero of the heavy particle corresponds to infinite time of the light one and the evolution in time of the system is completely neglected. In order to restore the time evolution, in [JZ] the formula is modified introducing by hand the internal dynamics of the heavy particle.

Our aim in this note is to discuss how the complete Joos and Zeh formula, i.e. modified taking into account the internal motion of the heavy particle, can be rigorously derived from the Schrödinger equation of the whole two-particle system.

In particular, using the result proved in [DFT], we shall write the asymptotic form of the wave function of the system approximating the true wave function in a specific scaling limit (involving the mass ratio and the strength of the interaction) with an explicit control of the error.

Furthermore we shall apply the result for the analysis of decoherence in a concrete example of quantum evolution.

More precisely, we shall consider an initial state with the heavy particle in a coherent superposition of two spatially separated wave packets with opposite momentum and the light one localized far on the left with a positive momentum.

Under precise assumptions on the relevant physical parameters (spreading and momentum of the light particle, effective range of the interaction, spreading and separation of the wave packets of the heavy particle) we shall derive an approximated form of the wave function of the system, describing the typical entangled state with the wave of the light particle splitted into a reflected and a transmitted part by each wave packet of the heavy particle.

As a consequence, the reduced density matrix for the heavy particle will show unperturbed diagonal terms and off-diagonal terms reduced by a factor Λ less than one, which gives a measure of the decoherence effect induced on the heavy particle.

Due to a phase shift, the waves reflected by the two wave packets are shown to be approximately orthogonal if the separation of the two wave packets is sufficiently large and then the factor Λ can be explicitly computed in terms of the transmission probability for the light particle subject to a δ -potential placed at a fixed position.

We want to emphasize that, at each step, the approximate formulas are obtained through rather elementary estimates with explicit control of the error. In this sense the model may be considered of pedagogical relevance for the analysis of the mechanism of decoherence.

Let us introduce the model. The hamiltonian of the two-particle system is given by

$$H = -\frac{\hbar^2}{2M}\Delta_R - \frac{\hbar^2}{2m}\Delta_r + \alpha_0\delta(r - R) \quad (2)$$

where M and m denote the mass of the heavy and the light particle respectively and $\alpha_0 > 0$ is the strength of the interaction.

The hamiltonian (2) is a well defined selfadjoint and positive operator in $L^2(\mathbb{R}^2)$ (see e.g. [AGH-KH]) and moreover it is a solvable model, in the sense that the corresponding generalized eigenfunctions and the unitary group can be explicitly computed ([S]).

We consider an initial state given in a product form

$$\psi_0(r, R) = \phi(R)\chi(r), \quad \phi, \chi \in \mathcal{S}(\mathbb{R}) \quad (3)$$

where $\mathcal{S}(\mathbb{R})$ is the Schwartz space, and we denote by $\epsilon = \frac{m}{M}$ the mass ratio which is the small parameter of the model.

We are interested in the asymptotic form of the solution of the corresponding Schrödinger equation

$$i\hbar \frac{\partial \psi_t^\epsilon}{\partial t} = H \psi_t^\epsilon \quad (4)$$

when $\epsilon \rightarrow 0$ and $\hbar$, M and the parameter

$$\alpha = \frac{m\alpha_0}{\hbar^2} \quad (5)$$

are kept fixed. Notice that α^{-1} is a length with the physical meaning of an effective range of the interaction.

Rescaling the time according to

$$\tau = \frac{\hbar}{M}t \quad (6)$$

the Schrödinger equation can be more conveniently written as

$$i \frac{\partial \psi_\tau^\epsilon}{\partial \tau} = -\frac{1}{2}\Delta_R \psi_\tau^\epsilon + \frac{1}{\epsilon} \left(-\frac{1}{2}\Delta_r + \alpha\delta(r - R) \right) \psi_\tau^\epsilon \quad (7)$$

From (7) it is clear that for $\epsilon \rightarrow 0$ the kinetic energy of the heavy particle can be considered as a small perturbation.

The situation is similar to the Born-Oppenheimer approximation, with the relevant difference that here the light particle is not in a bound state and so it cannot produce any effective potential for the heavy particle.

Then, for $\epsilon \rightarrow 0$, one should expect a scattering regime for the light particle in presence of the heavy one in a fixed position and a free motion for the heavy particle.

In fact, the asymptotics for $\epsilon \rightarrow 0$ of ψ_τ^ϵ is characterized in the following proposition ([DFT]).

Theorem 1. *For any initial state (3) and any $\tau > 0$ one has*

$$\|\psi_\tau^\epsilon - \psi_\tau^a\| < \left(\frac{A}{\tau} + B\right) \epsilon \quad (8)$$

where

$$\psi_\tau^a(r, R) = \frac{1}{\sqrt{i\epsilon^{-1}\tau}} e^{i\frac{\epsilon}{2\tau}r^2} \int dx e^{-i\tau H_0}(R-x)\phi(x) [(\Omega_+^x)^{-1}\chi]^\sim\left(\frac{r}{\epsilon^{-1}\tau}\right) \quad (9)$$

and $H_0 = -\frac{1}{2}\Delta$, the symbol $^\sim$ denotes the Fourier transform, Ω_+^x is the wave operator associated to the one-particle hamiltonian $H_x = -\frac{1}{2}\Delta + \alpha\delta(\cdot - x)$, for any $x \in \mathbb{R}$; moreover A, B are positive, time-independent constants whose dependence on the strength of the interaction and on the initial state are explicitly given.

We want to apply the result stated in theorem 1 to the analysis of decoherence in a concrete example of quantum evolution. In particular we consider an

initial state $\psi_0(R, r) = \phi(R)\chi(r)$ of the following form

$$\phi(R) = \frac{1}{\sqrt{2}} \left(f_\sigma^+(R) + f_\sigma^-(R) \right) \quad (10)$$

$$f_\sigma^\pm(R) = \frac{1}{\sqrt{\sigma}} f\left(\frac{R \pm R_0}{\sigma}\right) e^{\pm i P_0 R}, \quad \sigma, R_0, P_0 > 0, \quad R_0 > 2\sigma \quad (11)$$

$$\chi(r) = g_\delta(r) = \frac{1}{\sqrt{\delta}} g\left(\frac{r - r_0}{\delta}\right) e^{i q_0 r}, \quad \delta, q_0 > 0, \quad r_0 < -R_0 - \sigma - \delta \quad (12)$$

$$f, g \in C_0^\infty(-1, 1), \quad \|f\| = \|g\| = 1 \quad (13)$$

From (10),(11) one sees that the heavy particle is in a superposition state of two spatially separated wave packets, one localized in $R = -R_0$ with mean value of the momentum P_0 and the other localized in $R = R_0$ with mean value of the momentum $-P_0$.

From (12), the light particle is assumed localized around r_0 , on the left of the wave packet f_σ^+ , with positive mean momentum q_0 .

Moreover, to simplify the computation, f_σ^+ , f_σ^- , g_δ are chosen compactly supported and with disjoint supports.

We expect that in the time evolution of the above initial state, for $\epsilon \rightarrow 0$, the light particle will be partly reflected and partly transmitted by the two wave packets f_σ^+ , f_σ^- , which approximately act as fixed scattering centers.

We shall make precise this statement introducing suitable assumptions on the physical parameters of the system.

In order to formulate the result, we introduce the reflection and transmission coefficient associated to the one-particle hamiltonian H_x (see e.g. [AGH-KH])

$$\mathcal{R}_\alpha(k) = -\frac{\alpha}{\alpha - i|k|}, \quad \mathcal{T}_\alpha(k) = -\frac{i|k|}{\alpha - i|k|} \quad (14)$$

and define the transmitted and reflected part of the wave function of the light particle

$$g_{\mathcal{T}}(\epsilon^{-1}\tau, r) = \mathcal{T}_\alpha\left(\frac{|r|}{\epsilon^{-1}\tau}\right) \frac{e^{i\frac{r^2}{2\epsilon^{-1}\tau}}}{\sqrt{i\epsilon^{-1}\tau}} \tilde{g}_\delta\left(\frac{r}{\epsilon^{-1}\tau}\right) \quad (15)$$

$$g_{\mathcal{R}}^x(\epsilon^{-1}\tau, r) = \mathcal{R}_\alpha\left(\frac{|r|}{\epsilon^{-1}\tau}\right) e^{-2ix\frac{r}{\epsilon^{-1}\tau}} \frac{e^{i\frac{r^2}{2\epsilon^{-1}\tau}}}{\sqrt{i\epsilon^{-1}\tau}} \tilde{g}_\delta\left(-\frac{r}{\epsilon^{-1}\tau}\right) \quad (16)$$

Moreover, we shall assume

$$q_0 \gg \frac{1}{\delta}, \quad \frac{1}{\alpha} \gg \sigma \quad (17)$$

The first assumption in (17) simply means that the wave function of the light particle is well concentrated in momentum space around q_0 (notice that δ^{-1} is the order of magnitude of the spreading of the momentum).

The second assumption in (17) requires that the effective range of the interaction is much larger than the spreading in position of each wave packet of the heavy particle and this means that the interaction cannot "distinguish" two different points in the supports of f_σ^+ and f_σ^- .

Then

Proposition 2. *For the initial state (10), (11), (12), (13) and any $\tau > 0$ one has*

$$\|\psi_\tau^a - \psi_\tau^e\| < c \left[(\delta q_0)^{-n} + \sigma \alpha \right] \quad (18)$$

for any $n \in \mathbb{N}$, where c is a numerical constant depending on g and

$$\begin{aligned} \psi_\tau^e(r, R) &= \frac{1}{\sqrt{2}} \left(e^{-i\tau H_0} f_\sigma^+ \right) (R) \left[g_{\mathcal{T}}(\epsilon^{-1}\tau, r) + g_{\mathcal{R}}^{R_0}(\epsilon^{-1}\tau, r) \right] \\ &+ \frac{1}{\sqrt{2}} \left(e^{-i\tau H_0} f_\sigma^- \right) (R) \left[g_{\mathcal{T}}(\epsilon^{-1}\tau, r) + g_{\mathcal{R}}^{R_0}(\epsilon^{-1}\tau, r) \right] \end{aligned} \quad (19)$$

Proof. Exploiting the explicit expression of the generalized eigenfunctions of H_x (see e.g. [AGH-KH]), for $x \in \text{supp } f_\sigma^\pm$, one has

$$\begin{aligned}
[(\Omega_+^x)^{-1}g_\delta]^\sim(k) &= \frac{1}{\sqrt{2\pi}} \int dy g_\delta(y) \left(e^{-iky} + \mathcal{R}_\alpha(k) e^{-ikx} e^{i|k||x-y|} \right) \\
&= \mathcal{T}_\alpha(k) \tilde{g}_\delta(k) \theta_+(k) + \left[\tilde{g}_\delta(k) + e^{-2ikx} \mathcal{R}_\alpha(k) \tilde{g}_\delta(-k) \right] \theta_-(k)
\end{aligned} \tag{20}$$

where we have used the fact that $|x-y| = x-y$ for $x \in \text{supp } f_\sigma^\pm$, $y \in \text{supp } g_\delta$, the identity $1 + \mathcal{R}_\alpha = \mathcal{T}_\alpha$ and we have denoted by θ_+ , θ_- the characteristic functions of the positive and negative semiaxis respectively.

From (9),(10),(11),(12),(19) we can write

$$\begin{aligned}
&\psi_\tau^a(r, R) - \psi_\tau^e(r, R) \\
&= \frac{1}{\sqrt{2i\epsilon^{-1}\tau}} e^{i\frac{\epsilon}{2\tau}r^2} \int dx e^{-i\tau H_0}(R-x) (f_\sigma^+(x) + f_\sigma^-(x)) \mathcal{R}_\alpha\left(\frac{r}{\epsilon^{-1}\tau}\right) \left[\tilde{g}_\delta\left(\frac{r}{\epsilon^{-1}\tau}\right) \theta_-(r) \right. \\
&\quad \left. - e^{-2ix\frac{r}{\epsilon^{-1}\tau}} \tilde{g}_\delta\left(-\frac{r}{\epsilon^{-1}\tau}\right) \theta_+(r) \right] \\
&\quad + \frac{1}{\sqrt{2}} \int dx e^{-i\tau H_0}(R-x) f_\sigma^+(x) \left(g_{\mathcal{R}}^x\left(\frac{r}{\epsilon^{-1}\tau}\right) - g_{\mathcal{R}}^{-R_0}\left(\frac{r}{\epsilon^{-1}\tau}\right) \right) \\
&\quad + \frac{1}{\sqrt{2}} \int dx e^{-i\tau H_0}(R-x) f_\sigma^-(x) \left(g_{\mathcal{R}}^x\left(\frac{r}{\epsilon^{-1}\tau}\right) - g_{\mathcal{R}}^{R_0}\left(\frac{r}{\epsilon^{-1}\tau}\right) \right) \\
&\equiv (\zeta_1 + \zeta_2 + \zeta_3)(r, R)
\end{aligned} \tag{21}$$

Noticing that $\tilde{g}_\delta(k) = e^{i(q_0-k)r_0} \sqrt{\delta} \tilde{g}(\delta k - \delta q_0)$, one has

$$\begin{aligned}
\|\zeta_1\|^2 &= \frac{1}{2} \int dx |f_\sigma^+(x) + f_\sigma^-(x)|^2 \int dk |\mathcal{R}_\alpha(k)|^2 \left| \tilde{g}_\delta(k) \theta_-(k) - e^{-2ikx} \tilde{g}_\delta(-k) \theta_+(k) \right|^2 \\
&\leq \int dk \left(|\tilde{g}_\delta(k) \theta_-(k)|^2 + |\tilde{g}_\delta(-k) \theta_+(k)|^2 \right) \\
&= 2 \int_{\delta q_0}^\infty dz |\tilde{g}(z)|^2
\end{aligned} \tag{22}$$

which is $O((\delta q_0)^{-n})$ for any $n \in \mathbb{N}$. Concerning ζ_2 we have

$$\begin{aligned}
\|\zeta_2\|^2 &= \frac{1}{2} \int dx |f_\sigma^+(x)|^2 \int dk |\mathcal{R}_\alpha(k)|^2 |\tilde{g}_\delta(k)|^2 \left| e^{-2ikx} - e^{2ikR_0} \right|^2 \\
&\leq 2 \sup_{x \in (-R_0-\sigma, -R_0+\sigma)} |x + R_0|^2 \int dk k^2 |\mathcal{R}_\alpha(k)|^2 |\tilde{g}_\delta(k)|^2 \\
&\leq 2\sigma^2 \alpha^2
\end{aligned} \tag{23}$$

where we have used the inequality $k^2 |\mathcal{R}_\alpha(k)|^2 \leq \alpha^2$. The estimate for ζ_3 proceeds exactly in the same way concluding the proof. $\square$

Proposition 2 shows that, under the assumptions (17), the wave function (19) of the whole system takes the form of the typical entangled state emerging from an interaction of von Neumann type between a system and an apparatus. Here the role of the apparatus is played by the light particle which is partly transmitted and partly reflected by each wave packet of the heavy particle.

In this process the light particle keeps the information about the position of the heavy particle, which is encoded in $g_{\mathcal{R}}^{-R_0}$ and $g_{\mathcal{R}}^{R_0}$, i.e. in the reflected waves by f_{σ}^{+} and f_{σ}^{-} respectively.

As it should be expected in a one dimensional scattering process, such reflected waves have an identical spatial localization and they only differ for a phase factor.

In order to discuss the decoherence effect induced on the heavy particle one has to consider the density matrix of the whole system $\rho_{\tau}^{\epsilon}(r, R, r', R')$ and then one should compute the reduced density matrix for the heavy particle

$$\hat{\rho}_{\tau}^{\epsilon}(R, R') \equiv \int dr \rho_{\tau}^{\epsilon}(r, R, r, R') \equiv \int dr \psi_{\tau}^{\epsilon}(r, R) \overline{\psi_{\tau}^{\epsilon}}(r, R') \quad (24)$$

which defines a positive, trace-class operator in $L^2(\mathbb{R}^2)$ with $Tr \hat{\rho}_{\tau}^{\epsilon} = 1$.

We also introduce the approximate reduced density matrix

$$\begin{aligned} \hat{\rho}_{\tau}^e(R, R') &\equiv \int dr \rho_{\tau}^e(r, R, r, R') \equiv \int dr \psi_{\tau}^e(r, R) \overline{\psi_{\tau}^e}(r, R') \\ &= \frac{1}{2} \left(e^{-i\tau H_0} f_{\sigma}^{+} \right) (R) \left(e^{i\tau H_0} \overline{f_{\sigma}^{+}} \right) (R') + \frac{1}{2} \left(e^{-i\tau H_0} f_{\sigma}^{-} \right) (R) \left(e^{i\tau H_0} \overline{f_{\sigma}^{-}} \right) (R') \\ &\quad + \frac{\Lambda}{2} \left(e^{-i\tau H_0} f_{\sigma}^{+} \right) (R) \left(e^{i\tau H_0} \overline{f_{\sigma}^{-}} \right) (R') + \frac{\bar{\Lambda}}{2} \left(e^{-i\tau H_0} f_{\sigma}^{-} \right) (R) \left(e^{i\tau H_0} \overline{f_{\sigma}^{+}} \right) (R') \end{aligned} \quad (25)$$

where, in the last equality, we took into account that

$$\int dr |g_{\mathcal{T}}(\epsilon^{-1}\tau, r) + g_{\mathcal{R}}^{-R_0}(\epsilon^{-1}\tau, r)|^2 = \int dr |g_{\mathcal{T}}(\epsilon^{-1}\tau, r) + g_{\mathcal{R}}^{R_0}(\epsilon^{-1}\tau, r)|^2 = 1 \quad (26)$$

and we defined

$$\Lambda \equiv \int dr \left(g_{\mathcal{T}}(\epsilon^{-1}\tau, r) + g_{\mathcal{R}}^{-R_0}(\epsilon^{-1}\tau, r) \right) \left(\overline{g_{\mathcal{T}}}(\epsilon^{-1}\tau, r) + \overline{g_{\mathcal{R}}^{R_0}}(\epsilon^{-1}\tau, r) \right) \quad (27)$$

Notice that $|\Lambda| \leq 1$ and the case $\Lambda = 1$ can only occurs if $\alpha = 0$, i.e. when the interaction is absent.

Since L^2 -convergence of the wave function implies convergence of the corresponding density matrix in the trace-class norm, we conclude that $\hat{\rho}_{\tau}^e$ is a good approximation in the trace-class norm of $\hat{\rho}_{\tau}^{\epsilon}$ under the assumptions of theorem 1 and proposition 2.

For $\alpha = 0$ we see from (25), (27) that $\hat{\rho}_{\tau}^e$ reduces to the pure state describing the coherent superposition of the two wave packets evolving in time according to the free hamiltonian.

Notice that the last two terms in (25), usually called non diagonal terms, are responsible for the interference effects observed when the two wave packets have a non empty overlapping.

The occurrence of the interference makes the state highly non classical and it is the distinctive character of a quantum system with respect to a classical one.

If we switch on the interaction, i.e. for $\alpha > 0$, the effect of the light particle is to reduce the non diagonal terms by the factor Λ and this means that the interference effects for the heavy particle are correspondingly reduced.

In this sense one can say that there is a (partial) decoherence effect induced on the heavy particle which is completely characterized by the parameter Λ .

In the limit $\Lambda \rightarrow 0$, i.e. for $\alpha \rightarrow \infty$, the non diagonal terms and the interference effects are completely cancelled and then the state becomes a classical statistical mixture of the two pure states $e^{-i\tau H_0} f_\sigma^+$, $e^{-i\tau H_0} f_\sigma^-$.

We conclude this note showing that the parameter Λ can be approximated by a simpler expression under further conditions on the parameters.

First we notice that using the first assumption in (17) one easily gets

$$\Lambda = \int dr |g_T(\epsilon^{-1}\tau, r)|^2 + \int dr g_{\mathcal{R}}^{-R_0}(\epsilon^{-1}\tau, r) \overline{g_{\mathcal{R}}^{R_0}}(\epsilon^{-1}\tau, r) + O((\delta q_0)^{-n}) \quad (28)$$

Moreover we can show that for a large separation of the two wave packets the second integral in (28) becomes negligible due to the rapid oscillations of the integrand. More precisely we assume

$$d \gg \frac{1}{\alpha}, \quad d \gg \delta, \quad d \equiv 2R_0 \quad (29)$$

From (16), (14) we have

$$\begin{aligned} \mathcal{I} &\equiv \int dr g_{\mathcal{R}}^{-R_0}(\epsilon^{-1}\tau, r) \overline{g_{\mathcal{R}}^{R_0}}(\epsilon^{-1}\tau, r) \\ &= \int dk |\mathcal{R}_\alpha(k)|^2 |\tilde{g}_\delta(-k)|^2 e^{2idk} = \int dz \frac{\alpha^2}{\alpha^2 + (\delta^{-1}z - q_0)^2} |\tilde{g}(z)|^2 e^{2id(\delta^{-1}z - q_0)} \end{aligned} \quad (30)$$

Integrating by parts in (30) one obtains

$$\begin{aligned} |\mathcal{I}| &= \left| \int dz \frac{\alpha^2}{\alpha^2 + (\delta^{-1}z - q_0)^2} |\tilde{g}(z)|^2 \frac{1}{2id\delta^{-1}} \frac{d}{dz} e^{2id(\delta^{-1}z - q_0)} \right| \\ &\leq \frac{\delta}{2d} \int dz \left| \frac{d}{dz} \left(\frac{\alpha^2}{\alpha^2 + (\delta^{-1}z - q_0)^2} |\tilde{g}(z)|^2 \right) \right| \\ &\leq \frac{\delta}{d} \int dz \frac{\alpha^2 \delta^{-1} |\delta^{-1}z - q_0|}{[\alpha^2 + (\delta^{-1}z - q_0)^2]^2} |\tilde{g}(z)|^2 + \frac{\delta}{d} \int dz \frac{\alpha^2}{\alpha^2 + (\delta^{-1}z - q_0)^2} |\tilde{g}(z)| |\tilde{g}'(z)| \\ &\leq \frac{1}{d\alpha} + \frac{\delta}{d} \|\tilde{g}'\| \end{aligned} \quad (31)$$

where, in the last line, we have used the trivial estimates

$$\frac{\alpha^2 \delta^{-1} |\xi|}{(\alpha^2 + \xi^2)^2} \leq \frac{\alpha^2}{\alpha^2 + \xi^2} \frac{\delta^{-1} |\xi|}{\alpha^2 + \xi^2} \leq \frac{\delta^{-1} |\xi|}{\alpha^2 + \xi^2} \leq \frac{1}{\delta \alpha} \quad (32)$$

and the Schwartz inequality.

Using the estimate (31) in (28) we conclude that

$$\left| \Lambda - \int dr |g_{\mathcal{T}}(\epsilon^{-1} \tau, r)|^2 \right| \leq \frac{1}{d\alpha} + \frac{\delta}{d} \|\tilde{g}'\| + O((\delta q_0)^{-n}) \quad (33)$$

This means that, under the assumptions (17),(29), the decoherence effect is measured by the transmission probability of the light particle which is explicitly given by

$$\int dr |g_{\mathcal{T}}(\epsilon^{-1} \tau, r)|^2 = \int dk |\mathcal{T}_{\alpha}(k)|^2 |\tilde{g}_{\delta}(k)|^2 \quad (34)$$

The previous analysis of decoherence induced by scattering is clearly limited by the consideration of a two-body system.

For a more satisfactory treatment one should consider a model with N light particles scattered by the heavy one. The expected result is that the effect of the scattering events is cumulative and then the decoherence effect is increased at each step ([JZ]).

We observe that a rigorous proof of this fact would require good estimates for the wave operator in a case of many-body scattering problem and it is a non trivial open question.

References

- [AGH-KH] Albeverio S., Gesztesy F., Hoegh-Krohn R., Holden H., *Solvable Models in Quantum Mechanics*, Springer-Verlag, 1988.
- [BGJKS] Blanchard Ph., Giulini D., Joos E., Kiefer C., Stamatescu I.-O. eds., *Decoherence: Theoretical, Experimental and Conceptual Problems*, Lect. Notes in Phys. 538, Springer, 2000.
- [D] Dell'Antonio G., On Decoherence, preprint 2003.
- [DFT] Duerr D., Figari R., Teta A., Decoherence in a Two-Particle Model, mp arc 02-442, submitted to *J. Math. Phys.*.
- [GF] Gallis M.R., Fleming G.N., Environmental and Spontaneous Localization, *Phys. Rev.* **A42**, 38-48 (1990).
- [GJKKSZ] Giulini D., Joos E., Kiefer C., Kupsch J., Stamatescu I.-O., Zeh H.D., *Decoherence and the Appearance of a Classical World in Quantum Theory*, Springer, 1996.
- [JZ] Joos E., Zeh H.D., The Emergence of Classical Properties Through Interaction with the Environment, *Z. Phys.* **B59**, 223-243 (1985).

- [S] Schulman L.S., Application of the propagator for the delta function potential, in *Path Integrals from meV to MeV*, Gutzwiller M.C., Ioumata A., Klauder J.K., Streit L. eds., World Scientific, 1986, pp. 302-311.
- [T] Tegmark M., Apparent Wave Function Collapse Caused by Scattering, *Found. Phys. Lett.* **6**, 571-590 (1993).